

**Claims:**

1. (Cancelled).
2. (Presently Amended) A method of enhancing **the** activity of an active pharmaceutical substance, **wherein said active pharmaceutical substance is phenazepam**, upon administration to a subject **suffering from a condition or disorder treatable by said active pharmaceutical substance in need thereof**, said method comprising combining **a therapeutic dose of** said active pharmaceutical substance with a homeopathically activated form of said **active pharmaceutical** substance.
3. (Previously Presented) The method of claim 2, wherein said combining step is carried out prior to said administration.
4. (Previously Presented) The method of claim 2, wherein the ratio of said homeopathically activated form to said active pharmaceutical substance is from 1:1 to 1:100.
5. (New) A method of enhancing the activity of an active pharmaceutical substance, wherein said active pharmaceutical substance is diazepam, upon administration to a subject suffering from a condition or disorder treatable by said active pharmaceutical substance, said method comprising combining a therapeutic dose of said active pharmaceutical substance with a homeopathically activated form of said active pharmaceutical substance.
6. (New) The method of claim 5, wherein said combining step is carried out prior to said administration.
7. (New) The method of claim 5, wherein the ratio of said homeopathically activated form to said active pharmaceutical substance is from 1:1 to 1:100.
8. (New) A method of enhancing the activity of an active pharmaceutical substance, wherein said active pharmaceutical substance is hydrocortisone, upon administration to a subject suffering from a condition or disorder treatable by said active pharmaceutical substance, said method comprising combining a therapeutic dose of said active pharmaceutical substance with a homeopathically activated form of said active pharmaceutical substance.

9. (New) The method of claim 8, wherein said combining step is carried out prior to said administration.

10. (New) The method of claim 8, wherein the ratio of said homeopathically activated form to said active pharmaceutical substance is from 1:1 to 1:100.

11. (New) A method of enhancing the activity of an active pharmaceutical substance, wherein said active pharmaceutical substance is cyclophosphamide, upon administration to a subject suffering from a condition or disorder treatable by said active pharmaceutical substance, said method comprising combining a therapeutic dose of said active pharmaceutical substance with a homeopathically activated form of said active pharmaceutical substance.

12. (New) The method of claim 11, wherein said combining step is carried out prior to said administration.

13. (New) The method of claim 11, wherein the ratio of said homeopathically activated form to said active pharmaceutical substance is from 1:1 to 1:100.

14. (New) A method of enhancing the activity of an active pharmaceutical substance, wherein said active pharmaceutical substance is ethanol, upon administration to a subject suffering from a condition or disorder treatable by said active pharmaceutical substance, said method comprising combining a therapeutic dose of said active pharmaceutical substance with a homeopathically activated form of said active pharmaceutical substance.

15. (New) The method of claim 14, wherein said combining step is carried out prior to said administration.

16. (New) The method of claim 14, wherein the ratio of said homeopathically activated form to said active pharmaceutical substance is from 1:1 to 1:100.

17. (New) A method of enhancing the activity of an active pharmaceutical substance, wherein said active pharmaceutical substance is morphine, upon administration to a subject suffering from a condition or disorder treatable by said

active pharmaceutical substance, said method comprising combining a therapeutic dose of said active pharmaceutical substance with a homeopathically activated form of said active pharmaceutical substance.

18. (New) The method of claim 17, wherein said combining step is carried out prior to said administration.

19. (New) The method of claim 17, wherein the ratio of said homeopathically activated form to said active pharmaceutical substance is from 1:1 to 1:100.